

Litteraturliste: The Patient Specific Functional Scale

(Sist oppdatert 21. oktober 2011).

Chatman AB, Hyams SP, Neel JM, Binkley JM, Stratford PW, Schomberg A, et al. The Patient-Specific Functional Scale: measurement properties in patients with knee dysfunction. *Phys Ther.* 1997;**77**:820-9.

Cleland, J. A., J. M. Fritz, J. M. Whitman, and J. A. Palmer, 2006, The reliability and construct validity of the Neck Disability Index and patient specific functional scale in patients with cervical radiculopathy: *Spine (Phila Pa 1976.)*, v. 31, no. 5, p. 598-602.

Donnelly, C., and A. Carswell, 2002, Individualized outcome measures: a review of the literature: *Canadian Journal of Occupational Therapy. Revue Canadienne d Ergotherapie*, v. 69, no. 2, p. 84-94.

Gross DP, Battie MC, Asante AK. The Patient-Specific Functional Scale: validity in workers' compensation claimants. *Arch Phys Med Rehabil.* 2008;**89**:1294-9.

Hall AM, Maher CG, Latimer J, Ferreira ML, Costa LO. The patient-specific functional scale is more responsive than the Roland Morris disability questionnaire when activity limitation is low. *Eur Spine J.* 2010.

Hart, D. L., and M. W. Werneke, 2004, Re: Pengel LHM, Refshauge KM, Maher CG. Responsiveness of pain, disability, and physical impairment outcomes in patients with low back pain. *Spine.* 2004;**29**:879-83: *Spine (Phila Pa 1976.)*, v. 29, no. 21, p. 2475-2476.

Jolles BM, Buchbinder R, Beaton DE. A study compared nine patient-specific indices for musculoskeletal disorders. *J Clin Epidemiol.* 2005;**58**:791-801.

Maughan, E. F., and J. S. Lewis, 2010, Outcome measures in chronic low back pain: *Eur. Spine J.*, v. 19, no. 9, p. 1484-1494.

McMillan CR, Binhammer PA. Which outcome measure is the best? Evaluating responsiveness of the Disabilities of the Arm, Shoulder, and Hand Questionnaire, the Michigan Hand Questionnaire and the Patient-Specific Functional Scale following hand and wrist surgery. *Hand (N Y).* 2009;**4**:311-8.

Mills K, Blanch P, Dev P, Martin M, Vicenzino B. A randomised control trial of short term efficacy of in-shoe foot orthoses compared with a wait and see policy for anterior knee pain and the role of foot mobility. *Br J Sports Med.* 2011.

Moe RH, Uhlig T, Kjekken I, Hagen KB, Kvien TK, Grotle M. Multidisciplinary and multifaceted outpatient management of patients with osteoarthritis: protocol for a randomised, controlled trial. *BMC Musculoskelet Disord.* 2010;**11**:253.

Nourbakhsh MR, Fearon FJ. The effect of oscillating-energy manual therapy on lateral epicondylitis: a randomized, placebo-control, double-blinded study. *J Hand Ther.* 2008;**21**:4-13.

O'Shea SD, Taylor NF, Paratz JD. Measuring changes in activity limitation and participation restriction in people with COPD. *Int J Ther Rehab*. 2005;**12**:264-8.

Pengel LH, Refshauge KM, Maher CG. Responsiveness of pain, disability, and physical impairment outcomes in patients with low back pain. *Spine*. 2004;**29**:879-83.

Pietrobon R, Coeytaux RR, Carey TS, Richardson WJ, DeVellis RF. Standard scales for measurement of functional outcome for cervical pain or dysfunction: a systematic review. *Spine*. 2002;**27**:515-22.

Poulsen E, Christensen HW, Roos EM, Vach W, Overgaard S, Hartvigsen J. Non-surgical treatment of hip osteoarthritis. Hip school, with or without the addition of manual therapy, in comparison to a minimal control intervention: protocol for a three-armed randomized clinical trial. *BMC Musculoskelet Disord*. 2011;**12**:88.

Russell TG, Buttrum P, Wootton R, Jull GA. Internet-based outpatient telerehabilitation for patients following total knee arthroplasty: a randomized controlled trial. *J Bone Joint Surg Am*. 2011;**93**:113-20.

Saner J, Kool J, de Bie RA, Sieben JM, Luomajoki H. Movement control exercise versus general exercise to reduce disability in patients with low back pain and movement control impairment. A randomised controlled trial. *BMC Musculoskelet Disord*. 2011;**12**:207.

Spanjer J, Groothoff JW, Brouwer S. Instruments used to assess functional limitations in workers applying for disability benefit: a systematic review. *Disabil Rehabil*. 2011.

Stratford PW, Gill C, Westaway MD, Binkley JM. Assessing disability and change on individual patients: A report of a patient specific measure. *Physiotherapy Canada*. 1995;**47**:258-63.

Walker MJ. Manual physical therapy examination and intervention of a patient with radial wrist pain: a case report. *J Orthop Sports Phys Ther*. 2004;**34**:761-9.

Wang WT, Olson SL, Campbell AH, Hanten WP, Gleeson PB. Effectiveness of physical therapy for patients with neck pain: an individualized approach using a clinical decision-making algorithm. *Am J Phys Med Rehabil*. 2003;**82**:203-18.

Westaway MD, Stratford PW, Binkley JM. The patient-specific functional scale: validation of its use in persons with neck dysfunction. *J Orthop Sports Phys Ther*. 1998;**27**:331-8.

Young, I. A., J. A. Cleland, L. A. Michener, and C. Brown, 2010, Reliability, construct validity, and responsiveness of the neck disability index, patient-specific functional scale, and numeric pain rating scale in patients with cervical radiculopathy: *American Journal of Physical Medicine and Rehabilitation*, v. 89, no. 10, p. 831-839.